

Next generation long baseline experiments on the path to leptonic CP violation

P. Migliozzi and F. Terranova

- ✓ On-peak versus off-peak experiments
- ✓ JHF-SK in the "null result" scenario
- ✓ JHF-SK and CNGS synergies

Oscillation probability

Taylor expansion around $\alpha \equiv \Delta m_{21}^2 / \Delta m_{31}^2$ and $\sin^2 2\vartheta_{13}$ for constant matter density:

$$\begin{aligned}
 P(\nu_\mu \rightarrow \nu_e) &\simeq \sin^2 2\vartheta_{13} \sin^2 \vartheta_{23} \frac{\sin^2[(1-\hat{A})\Delta]}{(1-\hat{A})^2} && \mathcal{O}_1 \text{ (leading term)} \\
 &- \alpha \sin 2\vartheta_{13} \xi \sin \delta_{CP} \sin \Delta \frac{\sin[\hat{A}\Delta] \sin[(1-\hat{A})\Delta]}{\hat{A} (1-\hat{A})} && \mathcal{O}_2 \text{ } (\sim \sin \Delta) \\
 &+ \alpha \sin 2\vartheta_{13} \xi \cos \delta_{CP} \cos \Delta \frac{\sin[\hat{A}\Delta] \sin[(1-\hat{A})\Delta]}{\hat{A} (1-\hat{A})} && \mathcal{O}_3 \text{ } (\sim \cos \Delta) \\
 &+ \alpha^2 \cos^2 \vartheta_{23} \sin^2 2\vartheta_{12} \frac{\sin^2[\hat{A}\Delta]}{\hat{A}^2} && \mathcal{O}_4 \text{ (suppressed by } \alpha^2)
 \end{aligned}$$
$$\xi \equiv \cos \vartheta_{13} \sin 2\vartheta_{12} \sin 2\vartheta_{23} \sim \mathcal{O}(1) \qquad \hat{A} \equiv 2\sqrt{2} G_F n_e \frac{E}{\Delta m_{31}^2} \qquad \Delta \equiv \frac{\Delta m_{31}^2 L}{4E}$$


A.Cervera et al. Nucl.Phys. B579(2000)17 ; M. Freund, Phys.Rev. D64(2001)053003

Hierarchy of $O_1 \dots O_4$ terms

- On peak, "short" baseline experiments (JHF-SK) \Rightarrow dominance of O_1 and O_2 terms and low sensitivity to sign (Δm_{31}^2)

$$P(\nu_\mu \rightarrow \nu_e) \simeq \sin 2 \vartheta_{13} (\sin 2 \vartheta_{13} A_1 - \sin \delta \alpha A_2) \quad ; \quad A_1, A_2 \sim O(1)$$

- On peak, longer baseline experiments (NuMI-Off Axis) \Rightarrow dominance of O_1 and O_2 and higher dependence on sign(Δm_{31}^2)




Off-peak experiments (e.g. CNGS)

Leading term: signal rate suppressed $|(1 - \hat{A})\Delta| \ll 1$

$$\frac{\sin^2[(1 - \hat{A})\Delta]}{(1 - \hat{A})^2} \simeq \Delta^2$$

Matter effects cancel out at LO even if CNGS is an high energy beam

$$P(\nu_\mu \rightarrow \nu_e) \simeq \Delta^2 [\sin^2 2\vartheta_{13} A_1 - \sin \delta \sin 2\vartheta_{13} \alpha \Delta A_2 + \cos \delta \sin 2\vartheta_{13} \alpha A_3 + \alpha^2 A_4]$$




Dominance of O_1 and O_3 :

O_1 is CP and matter independent

O_3 is CP even and odd under

$\Delta m_{31}^2 \rightarrow -\Delta m_{31}^2$ transformation



Phase I \rightarrow Phase II strategy

Since the physics reach of High Intensity Superbeams (e.g. JHF–HK) and NuFact depends critically on the size of $\sin^2 2\vartheta_{13}$

Phase I experiments \Rightarrow high $\sin^2 2\vartheta_{13}$ sensitivity

signal \Rightarrow precision MNS physics at SB/NuFact

null result \Rightarrow discourage the SB/NuFact physics programme


Three ways to build a good phase I experiment:

- A "pure" $\sin^2 2\vartheta_{13}$ experiments (e.g. Reactors)
- An experiment sensitive to δ but able to disentangle δ – ϑ_{13} cancellation effects (JHF–SK + antineutrino runs)
- An experiment which has maximal ϑ_{13} sensitivity for maximal

~~CP~~

How looks a pure $\sin^2 2\vartheta_{13}$ experiment?



T.Nakaya @ v2002





JHF-SK limit imposing $\delta_{CP}=0 \Delta m^2 > 0$

What happens in real life?

JHF-SK has the wrong pattern for $\delta > 0$




CNGS has the wrong pattern for $\Delta m^2_{31} > 0$




What can we say on Phase II if we observe a null result in JHF-SK?

Assuming complete ignorance on δ_{CP} and using no other information to lift the $\theta_{13}-\delta_{CP}$ ambiguity...



P.Huber et al.
Nucl.Phys.B645
(2002) 3

Even worse for higher $\Delta m^2_{21}/\Delta m^2_{31}$



HLMA solution

JHF–SK should exploit its higher sensitivity even in case of null results along the line of:

T.Kajita et al. Phys.Lett. B528 (2002) 245
(anti- ν with JHF–SK)

H.Minakata et al. hep-ph/0301210
(ν with JHF–SK and anti- ν with NuMI OA... "hey guys, no kidding: YOU japanese will run with anti- ν first!")

Can we exploit the different δ_{CP} patterns even in case of positive signal at JHF?

NuMI OA re-tuned: see P.Huber et al. Nucl.Phys. B654 (2003) 3
What about CNGS?


Status of CNGS at the beginning of JHF-SK

After 3 years data taking at the CNGS

- Evidence for τ appearance at $>2\sigma$ (OPERA alone)
- $\text{Sin}^2 2\theta_{13} < 0.035$ @90% C.L. (a factor 4 better than CHOOZ) in the worst case
- Indication of ν_e appearance if $\theta_{13} > 7^\circ$

$\theta_{13} _{\text{true}}$	$\theta_{13} _{\text{min}}$	$\theta_{13} _{\text{max}}$	
1°	---	5.5°	---
2.5°	---	5.8°	---
5.0°	---	7.0°	---
7.5°	1.2°	11.4°	$(7.5+3.9-6.3)^\circ$
10°	5.6°	13.7°	$(10.+3.7-4.4)^\circ$



90% C.L. allowed region



Allowed regions for JHF-SK(5 years) + CNGS(8years)

$\theta_{13}=5^\circ$

$\theta_{13}=10^\circ$



$\Delta m^2 < 0$


$\Delta m^2 > 0$

$\Delta m^2 < 0$

$\Delta m^2 > 0$

What about an anti-neutrino run at the CNGS?

Here we assumed CNGS 3 years ν and 6 years anti- ν



Conclusions

- Off-peak beams like CNGS or a re-tuned NuMI-OffAxis explore peculiar regions of the $\nu_{\mu} \rightarrow \nu_{\tau}$ oscillation probability that could complement on-peak experiments
- Comparisons between JHF-SK and CNGS in case of null result give the most convincing evidence that the JHF-SK data taking should be better optimized to fully profit of its overwhelming physics potential
- If $\theta_{13} > 7^\circ$, after three years data taking CNGS could give a first indication of ν_e appearance. Moreover, the CNGS is an off-peak beam, therefore it has a different pattern from JHF-SK \Rightarrow they can be used synergically